

MILES TO GO

THE RESPONSE TO HIV
IN THE CARIBBEAN



Contents

| | |
|----------------------------|-----------|
| Caribbean | 2 |
| Bahamas | 10 |
| Barbados | 12 |
| Belize | 14 |
| Cuba | 16 |
| Dominican Republic | 18 |
| Guyana | 20 |
| Haiti | 22 |
| Jamaica | 24 |
| Suriname | 26 |
| Trinidad and Tobago | 28 |
| | |
| Annex on methods | 31 |

Caribbean

AT A GLANCE

1 Renewed commitment to combination prevention that is tailored to populations and locations with the greatest need is required to accelerate reductions in new HIV infections.


2 Particular emphasis must be placed on the knowledge and service access gaps facing young people, men and key populations in the Caribbean.

3 Stigma and discrimination and restrictive laws and policies hinder access to services for young people and some key populations.

4 There is a large gap in awareness of HIV status at the start of the HIV testing and treatment cascade.

5 Proven strategies for early diagnosis and treatment enrolment, retention and adherence must be applied and scaled up rapidly to achieve the 90–90–90 targets.

HIV prevalence among young men and women (aged 15–24 years) and the existence of education policies that guide delivery of life skills-based HIV and sexuality education in secondary schools, Caribbean, 2017




Source: UNAIDS 2018 estimates; 2017 National Commitments and Policy Instrument; United Nations Population Fund 2018 state of the art diagnosis of comprehensive sexuality education (CSE) implementation in the English- and Dutch-speaking Caribbean (unpublished).

Nearly 90% of new infections in the Caribbean in 2017 occurred in four countries—Cuba, Dominican Republic, Haiti and Jamaica—while 87% of deaths from AIDS-related illness occurred in Dominican Republic, Haiti and Jamaica.

Haiti alone accounts for nearly half of new HIV infections and deaths due to AIDS-related illness. It is one of the few countries in the region that does not provide comprehensive sexuality education in primary and secondary schools. Comprehensive sexuality education plays a central role in the preparation of adolescents and young people for a safe, productive and fulfilling life, and it is an important component of the HIV prevention package for young people.


EPIDEMIC TRANSITION MEASURES

Number of new HIV infections, Caribbean, 2000–2017




Source: UNAIDS 2018 estimates.

Number of AIDS-related deaths, Caribbean, 2000–2017



Source: UNAIDS 2018 estimates.


Incidence:prevalence ratio, Caribbean, 1990–2017



... Target value

Source: UNAIDS 2018 estimates.

Number of new HIV infections and deaths among the HIV population, Caribbean, 1990–2017




Source: UNAIDS 2018 estimates.

There has been moderate progress on both prevention and treatment in the Caribbean. The annual number of new HIV infections among adults in the Caribbean declined by about 18% from 2010 to 2017, from 19 000 [14 000–31 000] to 15 000 [11 000–26 000]. Deaths from AIDS-related illness declined by 23% (from 13 000 [9 300–22 000] to 10 000 [7 100–17 000]) over the same period. As a result, the region's incidence:prevalence ratio is edging towards the 0.03 epidemic transmission benchmark, reaching 0.05 [0.04–0.08] in 2017.


Gay men and other men who have sex with men accounted for nearly a quarter of new infections in 2017. Efforts to reach men and boys, and particularly gay men and other men who have sex with men, are constrained by health services insufficiently tailored to their needs and limited community-based services. In total, key populations and their sexual partners represented two thirds of new infections in the region.

Distribution of new HIV infections, by population group, Caribbean, 2017



† Individuals in this category did not report any HIV-related risk behaviour.
Source: UNAIDS special analysis, 2018.

Distribution of new HIV infections and AIDS-related deaths by country, Caribbean, 2017



Source: UNAIDS 2018 estimates.

LAWS AND POLICIES SCORECARD

| Country | Criminalization of transgender people | Criminalization of sex work | Criminalization of same-sex sexual acts | Drug use or possession for personal use an offence | Parental consent for adolescents to access HIV testing | Spousal consent for married women to access SRH services | Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | Laws or policies restricting the entry, stay and residence of people living with HIV | Mandatory HIV testing for marriage, work or residence permits or for certain groups |
|----------------------------------|---------------------------------------|-----------------------------|---|--|--|--|---|--|---|
| Antigua and Barbuda | Red | Red | Orange | Red | Red | Green | Green | Green | |
| Bahamas | Green | Red | Green | Red | Red | Red | Red | Red | |
| Barbados | Green | Red | Orange (c) | Red | Red | Green | Green | Green | |
| Belize | Grey | Red (a) | Orange (d) | Red (e) | Red (a) | Grey | Grey | Grey | |
| Cuba | Green | Green | Orange | Red | Orange | Green | Red | Red | |
| Dominica | Green | Red | Orange | Red | Red | Green | Green | Green | |
| Dominican Republic | Green | Green | Green | Red | Orange | Green | Red | Red | |
| Grenada | Grey | Red (f) | Orange (b) | Red (g) | Grey | Grey | Grey | Grey | |
| Guyana | Grey | Red (a) | Orange (b) | Red (h) | Red (a) | Grey | Grey | Green | |
| Haiti | Green | Green | Green | Red (i) | Orange (j) | Green | Green | Green | |
| Jamaica | Green | Red | Orange | Red | Orange | Green | Orange | Green | |
| Saint Kitts and Nevis | Grey | Grey | Orange (b) | Red (k) | Red | Green | Grey | Red | |
| Saint Lucia | Green | Red | Orange | Red (l) | Red (a) | Green | Red (m) | Green | |
| Saint Vincent and the Grenadines | Grey | Grey | Orange (b) | Red (n) | Grey | Grey | Grey | Grey | |
| Suriname | Green | Red | Green | Red | Orange | Green | Red | Red | |
| Trinidad and Tobago | Grey | Red (a) | Orange (b) | Red (o) | Orange (p) | Grey | Grey | Grey | |


| | | | | | | | | |
|--|--|---|--|--|--|--|--|--|
| <p>Red: Criminalized and/or prosecuted Green: Neither criminalized nor prosecuted Grey: Data not available</p> | <p>Red: Any criminalization or punitive regulation of sex work Green: Sex work is not subject to punitive regulations or is not criminalized Orange: Issue is determined/differs at subnational level Grey: Data not available</p> | <p>Red: Death penalty Orange: Imprisonment (14 years–life, up to 14 years) or no penalty specified Green: Laws penalizing same-sex sexual acts have been decriminalized or never existed, or no specific legislation Grey: Data not available</p> | <p>Red: Compulsory detention for drug offences Orange: Possession of drugs for personal use is specified as a criminal offence or drug use or consumption is a specific offence in law Green: Possession of drugs for personal use is specified as a non-criminal offence Grey: Data not available</p> | <p>Red: Yes, for adolescents younger than 18 Orange: Yes, for adolescents younger than 14 and 16 Green: Yes, for adolescents younger than 12 Blue: No Grey: Data not available</p> | <p>Red: Yes Green: No Grey: Data not available</p> | <p>Red: Yes Orange: No, but prosecutions exist based on general criminal laws Green: No Grey: Data not available</p> | <p>Red: Yes Green: No Grey: Data not available</p> | <p>Red: Yes Green: No Grey: Data not available</p> |
|--|--|---|--|--|--|--|--|--|

Note: Data on laws restricting the entry, stay and residence of people living with HIV are currently undergoing a global review that will involve country validation. An update is expected by the end of 2018.

Sources: National Commitment and Policy Instrument, 2017 and 2018; supplemented by additional sources where noted (see references at end of chapter).


STIGMA AND DISCRIMINATION

Percentage of men and women aged 15–49 years who would not buy vegetables from a shopkeeper living with HIV, Caribbean, most recent data, 2012–2016




*Female respondents only.
Source: Population-based surveys, 2012–2016.

Percentage of people living with HIV who experienced discrimination in health-care settings, Belize, 2013




Source: People Living with HIV Stigma Index survey, Belize, 2013.

Percentage of key populations who reported having avoided HIV testing in the past 12 months due to stigma and discrimination, Cuba, 2017



Source: Integrated biological and behavioural surveys, 2017.

Percentage of ever-married or partnered women aged 15–49 years who experienced physical and/or sexual violence by an intimate partner in the past 12 months, countries with available data, Caribbean, 2000–2017



Source: Population-based surveys, 2000–2017.


Large proportions of people in the Caribbean harbour misconceptions about HIV that fuel stigma and discrimination (1). In Jamaica, for example, 71% of people said they would not buy vegetables from a vendor who is living with HIV, as did 58% of people in Haiti and 49% of people in Dominican Republic (2).

There have been notable efforts to address the high levels of stigma and discrimination experienced by key populations in several Caribbean countries. A regional transgender coalition also has been established to coordinate advocacy around issues relating to human rights, social justice and HIV. Recent data from Cuba are encouraging: less than 1% of gay men and other men who have sex with men and about 2% of female sex workers reported avoiding taking an HIV test in the previous 12 months due to stigma and discrimination (3).

Intimate partner violence is a major concern. About one in six adult women surveyed in Dominican Republic and Haiti said they had recently experienced intimate partner violence (4). In Cuba, about 7% of adult women reported similar experiences.

COMBINATION HIV PREVENTION

Availability of pre-exposure prophylaxis (PrEP), Caribbean, 2018




Levels of condom use at last sex with a non-regular partner among young men (aged 15–24 years) ranged from 67% (Belize) to 79% (Jamaica); among young women (aged 15–24 years), the range was 49% (Dominican Republic) to 57% (Jamaica).

HIV prevalence among gay men and other men who have sex with men is particularly high in Trinidad and Tobago (32%), Bahamas (25%) and Haiti (13%). Among transgender people, it is highest in Cuba (20%) and among prisoners, it is highest in Dominica (29%).

The Bahamas and Barbados were the only countries providing pre-exposure prophylaxis (PrEP) through the public health system in 2018, although PrEP is available through private providers in Dominican Republic, Jamaica and Suriname. It is not yet available in Cuba, Dominica or Haiti.

Source: 2017 and 2018 National Commitments and Policy Instrument.


HIV prevalence among key populations, countries with available data, Caribbean, 2015–2017



Source: 2018 Global AIDS Monitoring.

HIV TESTING AND TREATMENT

HIV testing and treatment cascade, Caribbean, 2017



Source: UNAIDS special analysis, 2018; see annex on methods for more details.


Among the 310 000 [260 000–420 000] people living with HIV in the Caribbean at the end of 2017, 73% [53–95%] were aware of their HIV status, an increase from 66% [48–92%] in 2016. The gap to achieving the first 90 of the 90–90–90 targets in 2017 was 54 800 people living with HIV.

About 181 000 [159 000–188 000] people in the region were accessing antiretroviral therapy in 2017, or 57% [42–80%] of all people living with HIV. The gap to achieving the first and second 90s of the 90–90–90 targets in 2017 was 74 700 people living with HIV.

The estimated percentage of people living with HIV who achieved viral suppression increased from 37% [27–51%] in 2016 to 40% [30–56%] in 2017. The gap to achieving all three 90s in 2017 was the viral suppression of an additional 103 000 people living with HIV.

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

Cascade of services for preventing vertical transmission and transmission rate, Caribbean, 2017




Source: UNAIDS 2018 estimates; 2018 Global AIDS Monitoring.

Seven countries and island states in the Caribbean have been validated as having eliminated mother-to-child transmission of HIV: Anguilla, Antigua and Barbuda, Bermuda, the Cayman Islands, Cuba, Montserrat and Saint Kitts and Nevis. The rate of mother-to-child transmission (including breastfeeding) in the Caribbean in 2017 was 13.3% [10.7–15.6%], among the lowest in the world and significantly lower than the 18.7 [15.2– 22.0] rate in 2010. Antiretroviral treatment prophylaxis to prevent vertical transmission of HIV and to protect the woman's own health was coverage was 75% [58– >95%] in 2017, and almost half (48%) of HIV-exposed infants received an early infant diagnosis before eight weeks of age.

INVESTMENT

HIV resource availability by source, 2000–2017, and projected resource needs by 2020, Caribbean



*Estimates for low- and middle-income countries per 2015 World Bank income level classification. All figures are expressed in constant 2016 US dollars.

Source: UNAIDS 2018 resource availability and needs estimates.

The financial resources available for HIV responses in the Caribbean increased until 2011; since then, it has declined, largely due to scaled back international support. Between 2006 and 2017, the availability of domestic resources increased 123.7%, while international resources have decreased by 16%. In 2017, the United States President's Emergency Plan for AIDS Relief (PEPFAR) provided 57% of the total HIV resources in the region; the Global Fund to Fight AIDS, Tuberculosis and Malaria (Global Fund) provided 8%.

The response in Haiti, with the largest epidemic in the region, is more than 90% externally funded and extremely reliant on international support. A UNAIDS analysis shows that Haiti would be unable to absorb even a moderate cut in donor financing (see Chapter 7).

In total, approximately US\$ 315 million was available for the Caribbean's HIV programmes in 2017, considerably less than the US\$ 604 million needed to finance the achievement of Fast-Track programme coverage targets by 2020. Domestic funding for prevention programmes is low.

References

1. People Living with HIV Stigma Index surveys, 2013–2016.
2. Demographic and Health Surveys, 2012–2016.
3. Integrated biological and behavioural surveys, 2017.
4. Demographic and Health Survey, 2012.

Additional sources for the laws and policies scorecard

- a. Sexual Rights Initiative database [database]. Sexual Rights Initiative; c2016 (<http://sexualrightsdatabase.org/map/21/Adult%20sex%20work>).
- b. State-sponsored homophobia. A world survey of sexual orientation laws: criminalisation, protection and recognition. ILGA; 2017 (https://ilga.org/downloads/2017/ILGA_State_Sponsored_Homophobia_2017_WEB.pdf).
- c. Barbados. Sexual Offences Act, 1992. Section 9 (<http://www2.ohchr.org/english/bodies/hrc/docs/ngos/lgbti2.pdf>).
- d. Belize. Criminal Code. Revised edition (2000). Section 53 (https://www.oas.org/juridico/mla/en/blz/en_blz-int-text-cc.pdf).
- e. Belize. Misuse of Drugs Act, Chapter 103. Revised edition (2000). Section 7 (http://www.cicad.oas.org/fortalecimiento_institucional/legislations/PDF/BZ/misuse_of_drugs_act.pdf).
- f. Grenada. Criminal Code, Chapter 72A (76 of 1958). Section 137 (30) (<https://prostitution.procon.org/sourcefiles/GrenadaCriminalCode.pdf>).
- g. Grenada. Drug Abuse (Prevention and Control) Act (7 of 1992). Chapter 84A, Sec. 6 (<http://www.easterncaribbeanlaw.com/wp-content/uploads/2014/07/Microsoft-Word-Cap84A-Drug-Abuse-Prevention-and-Control-Act.doc.pdf>).
- h. Republic of Guyana. Narcotic Drugs and Psychotropic Substances (Control) Act (2 of 1988). Chapter 10:10, Sec. 4 (<http://dpi.gov.gy/wp-content/uploads/2016/06/Cap.-1010-Narcotic-Drugs-and-Psychotropic-Substances-Control.pdf>).
- i. Loi Relative au Contrôle et à la Répression du Trafic Illicite de la Drogue du 4 Octobre 2001. Article 49 (https://www.unodc.org/res/cld/document/hti/loi-2001_html/LOI_RELATIVE_AU_CONTROLE_ET_A_LA_REPRESSION_DU_TRAFIC_ILLICITE_DE_LA_DROGUE.pdf).
- j. Normes et Directives Nationales pour le Conseil et le Dépistage du VIH. Republic of Haiti; December 2014:29 (<https://mspp.gouv.ht/site/downloads/NORMES%20ET%20DIRECTIVES%20NATIONALES%208%20juin%202015.pdf>).
- k. St Christopher and Nevis. Drugs (Prevention and Abatement of the Misuse and Abuse of Drugs) Act. Revised edition. Sec. 6(1) (http://www.easterncaribbeanlaw.com/wp-content/uploads/2014/08/drugs_act.pdf).
- l. Saint Lucia. Drugs (Prevention of Misuse) Act. Revised Edition (2008). Chapter 3.02. Sec. 8(1) (<http://www.easterncaribbeanlaw.com/wp-content/uploads/2014/07/Drugs-Prevention-of-Misuse-Act-Cap.3.02.pdf>).
- m. Saint Lucia. Criminal Code. Revised Edition (2005). Chapter 3.01. Sec. 140 (<http://www.govt.lc/www/legislation/Criminal%20Code.pdf>).
- n. Saint Vincent and the Grenadines. Drugs (Prevention of Misuse) Act. Revised edition 1990. Chapter 219. Sec. 7(1) (http://www.cicad.oas.org/fortalecimiento_institucional/legislations/PDF/VC/drugs_act.pdf).
- o. Republic of Trinidad and Tobago. Dangerous Drugs Act (38 of 1991). Chapter 11:25. Sec. 5 (http://rgd.legalaffairs.gov.tt/laws2/alphabetical_list/lawspdfs/11.25.pdf).
- p. National HIV testing and counselling policy. Port of Spain: Ministry of Health, Trinidad and Tobago; 2006 (www.health.gov.tt/downloads/DownloadItem.aspx?id=258).

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|----------------------|----------------------|----------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | <500 [<500– <500] | <500 [<200– <500] | <200 [<200– <200] |
| New HIV infections (0–14) | ... [...–...] | ... [...–...] | ... [...–...] |
| New HIV infections (women, 15+) | <200 [<200– <200] | <100 [<100– <100] | <100 [<100– <100] |
| New HIV infections (men, 15+) | <200 [<200– <200] | <200 [<200– <200] | <100 [<100– <200] |
| HIV incidence per 1000 population | 0.93 [0.84–1.02] | 0.67 [0.57–0.77] | 0.38 [0.3–0.45] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | ... [...–...] | ... [...–...] | ... [...–...] |
| AIDS-related deaths (0–14) | ... [...–...] | ... [...–...] | ... [...–...] |
| AIDS-related deaths (women, 15+) | ... [...–...] | ... [...–...] | ... [...–...] |
| AIDS-related deaths (men, 15+) | ... [...–...] | ... [...–...] | ... [...–...] |
| People living with HIV | | | |
| People living with HIV (all ages) | 5300 [4600–6100] | 5700 [5000–6300] | 5300 [4600–6000] |
| People living with HIV (0–14) | ... [...–...] | ... [...–...] | ... [...–...] |
| People living with HIV (women, 15+) | 2400 [2100–2800] | 2600 [2400–3000] | 2300 [2100–2700] |
| People living with HIV (men, 15+) | 2800 [2300–3300] | 2900 [2500–3400] | 2800 [2400–3300] |

LAWS AND POLICIES

| | |
|---|--|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | Yes |
| Criminalization of sex work | Selling and buying sexual services is criminalized |
| Criminalization of same-sex sexual acts | Laws penalizing same-sex sexual acts have been decriminalized or never existed |
| Drug use or possession for personal use is an offence | Possession of drugs for personal use is specified as a criminal offence |
| Criminalization of transgender people | Neither criminalized nor prosecuted |
| Laws or policies restricting the entry, stay and residence of people living with HIV | No |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 18 years |
| Spousal consent for married women to access sexual and reproductive health services | Yes |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | Yes |

STIGMA AND DISCRIMINATION

Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent


VIOLENCE

Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months


EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2009 | US\$ 12 749 | US\$ 4 402 073 | ... | ... | US\$ 301 889 | US\$ 4 901 265 |


EPIDEMIC TRANSITION METRICS




Change in new HIV infections since 2010 = **-38%**



Change in AIDS-related deaths since 2010 =



Incidence:prevalence ratio = **0.03**




KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | ... | 2 085 | ... | ... | ... |
| HIV prevalence | ... | 25% | ... | ... | 2.2% |
| Know their HIV status | ... | 91% | ... | ... | ... |
| Antiretroviral therapy coverage | ... | ... | ... | ... | 71.4% |
| Condom use | ... | 63% | ... | ... | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV COMORBIDITIES

| | |
|---|------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 41 [27–59] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | 6.1% |
| Cervical cancer screening of women living with HIV | ... |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment | ... |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment | ... |

HIV TESTING AND TREATMENT CASCADE



| | | | |
|-----------------|--------------------|--------------------|--------------------|
| All ages | ...% [...-...%] | 57% [49–64%] | 39% [34–44%] |
| Children (0–14) | ...% [...-...%] | ...% [...-...%] | ...% [...-...%] |
| Women (15+) | ...% [...-...%] | 63% [56–71%] | 46% [40–52%] |
| Men (15+) | ...% [...-...%] | 51% [43–59%] | 34% [28–39%] |

| | |
|---|----|
| Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? | No |
|---|----|

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|--------------------|--------------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | ...% [...-...%] | ...% [...-...%] |
| Early infant diagnosis | ...% [...-...%] | ...% [...-...%] |

HIV PREVENTION

| | |
|--|----------------|
| Knowledge of HIV prevention among young people aged 15–24 years (2014) | |
| — Women | 2.8% |
| — Men | 5.8% |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) | |
| — Women | ... |
| — Men | ... |
| Women aged 15–49 years who have their demand for family planning satisfied by modern methods | ... |
| Men aged 15–49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | 9 |

Harm reduction

| | |
|---|-----|
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | No |
| — Safe injection rooms available (2016) | No |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|----------------------|----------------------|----------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | <200 [<200– <500] | <200 [<200– <500] | <200 [<200– <200] |
| New HIV infections (0–14) | ... [...–...] | ... [...–...] | ... [...–...] |
| New HIV infections (women, 15+) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| New HIV infections (men, 15+) | <200 [<200– <200] | <200 [<100– <200] | <200 [<100– <200] |
| HIV incidence per 1000 population | 0.71 [0.62–0.78] | 0.64 [0.55–0.74] | 0.57 [0.47–0.68] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (0–14) | ... [...–...] | ... [...–...] | ... [...–...] |
| AIDS-related deaths (women, 15+) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (men, 15+) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| People living with HIV | | | |
| People living with HIV (all ages) | 1300 [1100–1400] | 2000 [1800–2300] | 2700 [2400–3000] |
| People living with HIV (0–14) | ... [...–...] | ... [...–...] | ... [...–...] |
| People living with HIV (women, 15+) | <500 [<500– <500] | 600 [540–680] | 860 [770–950] |
| People living with HIV (men, 15+) | 900 [780–1000] | 1400 [1200–1600] | 1800 [1600–2100] |

LAWS AND POLICIES

| | |
|---|--|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | No |
| Criminalization of sex work | Selling and buying sexual services is criminalized |
| Criminalization of same-sex sexual acts | Yes, imprisonment (up to 14 years) |
| Drug use or possession for personal use is an offence | Drug use or consumption is a specific offence in law |
| Criminalization of transgender people | Neither criminalized nor prosecuted |
| Laws or policies restricting the entry, stay and residence of people living with HIV | No |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 18 years |
| Spousal consent for married women to access sexual and reproductive health services | No |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | No |

STIGMA AND DISCRIMINATION

Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV **2012**
20.7*

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent

VIOLENCE


Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months

*Female respondents only


EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|-----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2013 | ... | US\$ 5 478 563 | US\$ 235 742 | ... | US\$ 10 256 837 | US\$ 15 971 143 |


EPIDEMIC TRANSITION METRICS




Change in new HIV infections since 2010 = -10%



Change in AIDS-related deaths since 2010 = 178%



Incidence:prevalence ratio = 0.06




KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | ... | 2 618 | ... | ... | ... |
| HIV prevalence | 0% | 2.8% | ... | ... | 1.3% |
| Know their HIV status | ... | ... | ... | ... | ... |
| Antiretroviral therapy coverage | ... | 35.3% | ... | ... | 1.4% |
| Condom use | ... | 58.2% | ... | ... | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV COMORBIDITIES

| | |
|---|---------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 3 [3-4] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | 0% |
| Women who tested positive for HIV among those screened for cervical cancer (programme data) (2017) | 2.3% |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment (2017) | 0% |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment (2017) | 0% |

HIV TESTING AND TREATMENT CASCADE



| | ... | 49% [43-55%] | ... |
|-----------------|------------|--------------|--------------|
| All ages | [...-...%] | | [...-...%] |
| Children (0-14) | [...-...%] | | [...-...%] |
| Women (15+) | [...-...%] | 65% [59-73%] | 48% [43-53%] |
| Men (15+) | [...-...%] | 42% [36-48%] | 30% [25-34%] |

| | |
|---|----|
| Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? | No |
|---|----|

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|-----------------|-----------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | ...% [...-...%] | ...% [...-...%] |
| Early infant diagnosis | ...% [...-...%] | ...% [...-...%] |

HIV PREVENTION

| | |
|--|----------------|
| Knowledge of HIV prevention among young people aged 15-24 years (2014) | |
| — Women | 47.6% |
| — Men | 44.8% |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) (2017) | |
| — Women | 19.7% |
| — Men | 42.1% |
| Women aged 15-49 years who have their demand for family planning satisfied by modern methods | ... |
| Men aged 15-49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | 1 |

Harm reduction

| | |
|---|-----|
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | ... |
| — Safe injection rooms available (2016) | ... |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|-----------------------|-----------------------|-----------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | <500 [<500–4600] | <500 [<500–4700] | <500 [<500–4900] |
| New HIV infections (0–14) | <100 [<100–660] | <100 [<100–900] | <100 [<100–580] |
| New HIV infections (women, 15+) | <200 [<100–1900] | <200 [<200–2000] | <200 [<200–2000] |
| New HIV infections (men, 15+) | <200 [<200–2200] | <200 [<200–2200] | <200 [<200–2000] |
| HIV incidence per 1000 population | 0.9 [0.85–21.29] | 0.88 [0.83–20.51] | 0.93 [0.87–19.92] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | <100 [<100–2200] | <100 [<100–2500] | <200 [<200–3700] |
| AIDS-related deaths (0–14) | <100 [<100– <500] | <100 [<100–830] | <100 [<100–820] |
| AIDS-related deaths (women, 15+) | <100 [<100–830] | <100 [<100–940] | <100 [<100–1600] |
| AIDS-related deaths (men, 15+) | <100 [<100–1000] | <100 [<100–1300] | <100 [<100–1800] |
| People living with HIV | | | |
| People living with HIV (all ages) | 2200 [2000–45 000] | 3100 [2800–52 000] | 4500 [4000–58 000] |
| People living with HIV (0–14) | <100 [<100–3700] | <100 [<100–3200] | <200 [<100–3600] |
| People living with HIV (women, 15+) | 980 [880–20 000] | 1400 [1300–24 000] | 2000 [1900–27 000] |
| People living with HIV (men, 15+) | 1200 [1000–22 000] | 1600 [1400–25 000] | 2300 [2100–29 000] |

LAWS AND POLICIES

| | |
|---|---|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | ... |
| Criminalization of sex work | Partial criminalization of sex work |
| Criminalization of same-sex sexual acts | Yes, imprisonment (up to 14 years) |
| Drug use or possession for personal use is an offence | Possession of drugs for personal use is specified as a criminal offence |
| Criminalization of transgender people | ... |
| Laws or policies restricting the entry, stay and residence of people living with HIV | ... |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 18 years |
| Spousal consent for married women to access sexual and reproductive health services | ... |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | ... |

STIGMA AND DISCRIMINATION

| | 2006 | 2016 |
|---|-------|-------------|
| Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV | 42.1* | 31.9 |
| Percentage of people living with HIV denied health services because of their HIV status in the last 12 months | | 2013 6.6 |
| Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent | | 2013 9.2 |

VIOLENCE


Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months

*Female respondents only

EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2012 | US\$ 118 050 | US\$ 832 345 | US\$ 555 482 | US\$ 517 357 | US\$ 640 356 | US\$ 2 836 772 |

EPIDEMIC TRANSITION METRICS




KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|--|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | ... | ... | ... | ... | 1 415 |
| HIV prevalence | ... | ... | ... | ... | 5.8% |
| Know their HIV status | ... | 44.3% | ... | ... | ... |
| Antiretroviral therapy coverage | ... | ... | ... | ... | ... |
| Condom use | ... | 81.4% | ... | 85.7% | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV COMORBIDITIES

| | |
|--|------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 40 [26–57] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | 4% |
| Cervical cancer screening of women living with HIV | ... |
| Proportion of people coinfectd with HIV and hepatitis B virus receiving combined treatment | ... |
| Proportion of people coinfectd with HIV and hepatitis C virus starting hepatitis C treatment | ... |

HIV TESTING AND TREATMENT CASCADE



| | ...% | 31% | ...% |
|------------------------|------------|------------|------------|
| All ages | [...-...%] | [28- >95%] | [...-...%] |
| Children (0–14) | [...-...%] | [52- >95%] | [...-...%] |
| Women (15+) | [...-...%] | [29- >95%] | [...-...%] |
| Men (15+) | [...-...%] | [25- >95%] | [...-...%] |

| | |
|--|-----|
| Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? | ... |
|--|-----|

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|--|----------------|----------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | 74% [63- >95%] | 41% [37- >95%] |
| Early infant diagnosis | 75% [3–88%] | 36% [2–40%] |

HIV PREVENTION

| | |
|---|----------------|
| Knowledge of HIV prevention among young people aged 15–24 years (2016) | |
| — Women | 41.4% |
| — Men | 45% |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) | |
| — Women | ... |
| — Men | ... |
| Women aged 15–49 years who have their demand for family planning satisfied by modern methods | ... |
| Men aged 15–49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | ... |
| Harm reduction | |
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | ... |
| — Safe injection rooms available (2016) | ... |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|-----------------------|---------------------------|---------------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | 1400 [1200–1600] | 2200 [1800–2500] | 1800 [1500–2200] |
| New HIV infections (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| New HIV infections (women, 15+) | <500 [<500– <500] | 620 [540–710] | 520 [<500–620] |
| New HIV infections (men, 15+) | 1000 [820–1200] | 1500 [1200–1800] | 1300 [990–1600] |
| HIV incidence per 1000 population | 0.13 [0.11–0.14] | 0.2 [0.17–0.22] | 0.17 [0.14–0.19] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | <200 [<100– <200] | <500 [<200– <500] | <500 [<500– <500] |
| AIDS-related deaths (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (women, 15+) | <100 [<100– <100] | <100 [<100– <200] | <200 [<200– <200] |
| AIDS-related deaths (men, 15+) | <100 [<100– <200] | <200 [<100– <500] | <200 [<100– <500] |
| People living with HIV | | | |
| People living with HIV (all ages) | 9000 [8000–10 000] | 17 000 [15 000–19 000] | 30 000 [26 000–33 000] |
| People living with HIV (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| People living with HIV (women, 15+) | 2500 [2200–2800] | 4700 [4200–5300] | 8000 [7100–9000] |
| People living with HIV (men, 15+) | 6500 [5600–7500] | 12 000 [11 000–14 000] | 21 000 [19 000–25 000] |

LAWS AND POLICIES

| | |
|---|---|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | Yes |
| Criminalization of sex work | Sex work is not subject to punitive regulations or is not criminalized |
| Criminalization of same-sex sexual acts | No penalty specified |
| Drug use or possession for personal use is an offence | Possession of drugs for personal use is specified as a criminal offence |
| Criminalization of transgender people | Neither criminalized nor prosecuted |
| Laws or policies restricting the entry, stay and residence of people living with HIV | Yes |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 14 years |
| Spousal consent for married women to access sexual and reproductive health services | No |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | Yes |

STIGMA AND DISCRIMINATION

| | 2006 | 2014 |
|---|-------|------|
| Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV | 22.4* | 16.4 |

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent

VIOLENCE


| | | |
|---|------|-----|
| Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months | 2017 | 6.8 |
|---|------|-----|

*Female respondents only


EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|-----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2013 | ... | US\$ 63 807 290 | ... | US\$ 5 927 082 | ... | US\$ 69 734 372 |


EPIDEMIC TRANSITION METRICS




Change in new HIV infections since 2010 = **-15%**



Change in AIDS-related deaths since 2010 = **31%**




Incidence:prevalence ratio = **0.06**



KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | 82 451 | 279 249 | ... | 3 700 | ... |
| HIV prevalence | 2.8% | 5.6% | ... | 19.7% | ... |
| Know their HIV status | ... | ... | ... | ... | ... |
| Antiretroviral therapy coverage | 86.7% | 83.4% | ... | 86.8% | ... |
| Condom use | 78.1% | 63.9% | ... | 65.8% | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | 1.9% | 0.7% | ... | ... | ... |

HIV TESTING AND TREATMENT CASCADE



| | | | |
|-----------------|------------------|-----------------|-----------------|
| All ages | 80% [70–90%] | 66% [58–75%] | 43% [38–49%] |
| Children (0–14) | 66% [58–75%] | 66% [58–75%] | 43% [38–49%] |
| Women (15+) | 57% [50–65%] | 49% [44–56%] | 33% [29–38%] |
| Men (15+) | 88% [76–>95%] | 72% [62–83%] | 47% [40–54%] |

Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? **Yes**

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|-------------------|--------------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | >95% [87–>95%] | >95% [93–>95%] |
| Early infant diagnosis | >95% [85–>95%] | >95% [>95–>95%] |

HIV COMORBIDITIES

| | |
|---|---------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 69 [44–98] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | 0.3% |
| Cervical cancer screening of women living with HIV | ... |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment (2017) | 83.8% |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment (2017) | 18.9% |

HIV PREVENTION

| | |
|--|----------------|
| Knowledge of HIV prevention among young people aged 15–24 years (2014) | |
| — Women | 60.9% |
| — Men | 58.6% |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) (2017) | |
| — Women | 75.5% |
| — Men | 80.2% |
| Women aged 15–49 years who have their demand for family planning satisfied by modern methods | ... |
| Men aged 15–49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | ... |

Harm reduction

| | |
|---|-----|
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | No |
| — Safe injection rooms available (2016) | No |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|----------------------------|----------------------------|---------------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | 5900 [4500–7800] | 2700 [1900–3800] | 2400 [1200–4900] |
| New HIV infections (0–14) | <500 [<500–560] | <200 [<100– <500] | <100 [<100– <200] |
| New HIV infections (women, 15+) | 2800 [2200–3700] | 1300 [930–1900] | 1100 [560–2300] |
| New HIV infections (men, 15+) | 2700 [2000–3700] | 1300 [860–1800] | 1200 [640–2700] |
| HIV incidence per 1000 population | 0.65 [0.5–0.87] | 0.28 [0.19–0.4] | 0.22 [0.12–0.46] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | 6500 [4500–9500] | 5000 [3500–7000] | 2600 [1700–3800] |
| AIDS-related deaths (0–14) | <500 [<500–520] | <200 [<100– <200] | <100 [<100– <100] |
| AIDS-related deaths (women, 15+) | 3000 [2100–4100] | 2100 [1500–3000] | 1200 [750–1800] |
| AIDS-related deaths (men, 15+) | 3200 [2200–4800] | 2700 [1900–3700] | 1300 [850–2000] |
| People living with HIV | | | |
| People living with HIV (all ages) | 99 000 [77 000–130 000] | 83 000 [62 000–100 000] | 67 000 [52 000–85 000] |
| People living with HIV (0–14) | 2500 [1800–3500] | 2000 [1500–2800] | 1400 [1000–1800] |
| People living with HIV (women, 15+) | 43 000 [33 000–56 000] | 38 000 [30 000–49 000] | 33 000 [26 000–42 000] |
| People living with HIV (men, 15+) | 54 000 [41 000–70 000] | 43 000 [32 000–55 000] | 33 000 [25 000–43 000] |

LAWS AND POLICIES

| | |
|---|---|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | Yes |
| Criminalization of sex work | Sex work is not subject to punitive regulations or is not criminalized |
| Criminalization of same-sex sexual acts | No specific legislation |
| Drug use or possession for personal use is an offence | Possession of drugs for personal use is specified as a criminal offence |
| Criminalization of transgender people | Neither criminalized nor prosecuted |
| Laws or policies restricting the entry, stay and residence of people living with HIV | Yes |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 16 years |
| Spousal consent for married women to access sexual and reproductive health services | No |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | Yes |

STIGMA AND DISCRIMINATION

| | 2007 | 2013 |
|---|------|------|
| Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV | 54.1 | 49.3 |

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent


VIOLENCE

| | 2007 | 2013 |
|---|------|------|
| Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months | 11.7 | 16 |


EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|-----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2017 | ... | US\$ 14 743 318 | ... | US\$ 6 230 084 | ... | US\$ 20 973 402 |


EPIDEMIC TRANSITION METRICS




Change in new HIV infections since 2010 = **-14%**



Change in AIDS-related deaths since 2010 = **-48%**



Incidence:prevalence ratio = **0.04**




KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | ... | 130 571 | ... | 9 327 | 19 994 |
| HIV prevalence | ... | ... | ... | ... | ... |
| Know their HIV status | ... | ... | ... | ... | ... |
| Antiretroviral therapy coverage | ... | ... | ... | ... | ... |
| Condom use | ... | ... | ... | ... | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV COMORBIDITIES

| | |
|---|---------------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 1500 [1100–2000] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | 21.1% |
| Cervical cancer screening of women living with HIV (survey data) (2013) | 81% |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment | ... |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment | ... |

HIV TESTING AND TREATMENT CASCADE



| | | | |
|-----------------|------------------|-----------------|--------------------|
| All ages | 77% [60–>95%] | 52% [41–66%] | 43% [34–55%] |
| Children (0–14) | 80% [59–>95%] | 58% [43–77%] | ...% [...–...%] |
| Women (15+) | 83% [66–>95%] | 56% [44–71%] | ...% [...–...%] |
| Men (15+) | 70% [53–92%] | 48% [37–63%] | ...% [...–...%] |

| | |
|---|----|
| Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? | No |
|---|----|

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|-----------------|-------------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | 59% [46–79%] | >95% [74–>95%] |
| Early infant diagnosis | 10% [8–13%] | 80% [62–>95%] |

HIV PREVENTION

| | |
|---|----------------|
| Knowledge of HIV prevention among young people aged 15–24 years (2014) | |
| — Women | 46.4% |
| — Men | ... |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) | |
| — Women | ... |
| — Men | ... |
| Women aged 15–49 years who have their demand for family planning satisfied by modern methods (2017) | 82.7% |
| Men aged 15–49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | ... |

| | |
|---|------|
| Harm reduction | |
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy (2017) | 7.8% |
| — Naloxone available (2016) | No |
| — Safe injection rooms available (2016) | No |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|----------------------|----------------------|----------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | 690 [590–850] | <500 [<500–610] | <500 [<500–580] |
| New HIV infections (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| New HIV infections (women, 15+) | <500 [<500– <500] | <200 [<200– <500] | <200 [<200– <500] |
| New HIV infections (men, 15+) | <500 [<500– <500] | <500 [<500– <500] | <500 [<200– <500] |
| HIV incidence per 1000 population | 1 [0.86–1.23] | 0.7 [0.55–0.88] | 0.62 [0.47–0.82] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | <200 [<100– <200] | <100 [<100– <200] | <200 [<200– <200] |
| AIDS-related deaths (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (women, 15+) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (men, 15+) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <200] |
| People living with HIV | | | |
| People living with HIV (all ages) | 5100 [4400–5700] | 6700 [6100–7400] | 8200 [7600–9200] |
| People living with HIV (0–14) | <200 [<200– <200] | <500 [<200– <500] | <500 [<200– <500] |
| People living with HIV (women, 15+) | 2600 [2300–3000] | 3100 [2900–3500] | 3800 [3400–4300] |
| People living with HIV (men, 15+) | 2300 [2000–2700] | 3300 [3000–3800] | 4300 [3800–4800] |

LAWS AND POLICIES

| | |
|---|---|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | ... |
| Criminalization of sex work | Buying sexual services is criminalized |
| Criminalization of same-sex sexual acts | Yes, imprisonment (up to 14 years) |
| Drug use or possession for personal use is an offence | Possession of drugs for personal use is specified as a criminal offence |
| Criminalization of transgender people | ... |
| Laws or policies restricting the entry, stay and residence of people living with HIV | ... |
| Parental consent for adolescents to access HIV testing | No |
| Spousal consent for married women to access sexual and reproductive health services | ... |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | No |

STIGMA AND DISCRIMINATION

| | 2005 | 2014 |
|---|------|------|
| Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV | 50.8 | 29.4 |

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent


VIOLENCE

Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months


EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|-----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2012 | US\$ 1 142 925 | US\$ 3 347 064 | ... | ... | US\$ 1 000 000 | US\$ 21 921 471 |


EPIDEMIC TRANSITION METRICS




Change in new HIV infections since 2010 = **-10%**



Change in AIDS-related deaths since 2010 = **64%**



Incidence:prevalence ratio = **0.05**




KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | 5 256 | 3 327 | ... | ... | ... |
| HIV prevalence | 6.1% | 4.9% | ... | 8.4% | 1.72% |
| Know their HIV status | ... | ... | ... | ... | ... |
| Antiretroviral therapy coverage | ... | ... | ... | ... | ... |
| Condom use | 75.7% | 64.4% | ... | ... | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV COMORBIDITIES

| | |
|---|------------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 170 [110–250] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | 4.7% |
| Cervical cancer screening of women living with HIV | ... |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment | ... |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment | ... |

HIV TESTING AND TREATMENT CASCADE



| | ... | 64% | ... |
|-----------------|------------|----------|------------|
| All ages | [...-...%] | [59–72%] | [...-...%] |
| Children (0–14) | [...-...%] | [53–71%] | [...-...%] |
| Women (15+) | [...-...%] | [68–85%] | [...-...%] |
| Men (15+) | [...-...%] | [49–62%] | [...-...%] |

| | |
|---|----|
| Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? | No |
|---|----|

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|-----------------|-----------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | 60% [53–66%] | 64% [55–71%] |
| Early infant diagnosis | 27% [24–30%] | 53% [47–61%] |

HIV PREVENTION

| | |
|--|----------------|
| Knowledge of HIV prevention among young people aged 15–24 years (2014) | |
| — Women | 51.5% |
| — Men | 40% |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) | |
| — Women | ... |
| — Men | ... |
| Women aged 15–49 years who have their demand for family planning satisfied by modern methods | ... |
| Men aged 15–49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | ... |

Harm reduction

| | |
|---|-----|
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | ... |
| — Safe injection rooms available (2016) | ... |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|--|-----------------------------|------------------------------|------------------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | 10 000 [7800–14 000] | 9800 [7500–13 000] | 7600 [5800–10 000] |
| New HIV infections (0–14) | 1500 [980–2100] | 1300 [850–1800] | 950 [640–1400] |
| New HIV infections (women, 15+) | 4500 [3400–6000] | 4400 [3300–5800] | 3400 [2600–4500] |
| New HIV infections (men, 15+) | 4300 [3200–6000] | 4200 [3100–5900] | 3200 [2400–4600] |
| HIV incidence per 1000 population | 1.2 [0.89–1.57] | 1.05 [0.78–1.38] | 0.73 [0.54–0.95] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | 7800 [5800–11 000] | 5300 [4000–7700] | 4700 [3600–6900] |
| AIDS-related deaths (0–14) | 1100 [660–1600] | 760 [<500–1100] | 600 [<500–910] |
| AIDS-related deaths (women, 15+) | 3200 [2300–4800] | 2100 [1600–3200] | 1800 [1300–2700] |
| AIDS-related deaths (men, 15+) | 3500 [2600–5200] | 2400 [1800–3600] | 2400 [1800–3500] |
| People living with HIV | | | |
| People living with HIV (all ages) | 110 000 [99 000–140 000] | 130 000 [110 000–160 000] | 150 000 [130 000–180 000] |
| People living with HIV (0–14) | 7400 [5900–9700] | 8000 [6300–11 000] | 7600 [5900–9800] |
| People living with HIV (women, 15+) | 56 000 [49 000–70 000] | 66 000 [56 000–80 000] | 78 000 [67 000–95 000] |
| People living with HIV (men, 15+) | 50 000 [43 000–63 000] | 56 000 [48 000–69 000] | 62 000 [53 000–77 000] |

LAWS AND POLICIES

| | |
|--|---|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | No |
| Criminalization of sex work | Sex work is not subject to punitive regulations or is not criminalized |
| Criminalization of same-sex sexual acts | No specific legislation |
| Drug use or possession for personal use is an offence | Possession of drugs for personal use is specified as a criminal offence |
| Criminalization of transgender people | Neither criminalized nor prosecuted |
| Laws or policies restricting the entry, stay and residence of people living with HIV | No |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 16 years |
| Spousal consent for married women to access sexual and reproductive health services | No |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | No |

STIGMA AND DISCRIMINATION

| | 2006 | 2012 |
|--|------|------|
| Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV | 68.3 | 57.7 |

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent


VIOLENCE

| | 2006 | 2012 |
|--|------|------|
| Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months | 17 | 14.9 |


EXPENDITURES

| | Financing sources | | | | | Total |
|------------------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|------------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2016 | ... | US\$ 3 968 260 | ... | ... | ... | US\$ 111 563 836 |


EPIDEMIC TRANSITION METRICS




Change in new HIV infections since 2010 = **-23%**



Change in AIDS-related deaths since 2010 = **-10%**



Incidence:prevalence ratio = **0.05**




KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | 70 302 | 30 853 | ... | ... | ... |
| HIV prevalence | ... | ... | ... | ... | 2.7% |
| Know their HIV status | ... | 69.5% | ... | ... | ... |
| Antiretroviral therapy coverage | ... | ... | ... | ... | ... |
| Condom use | 89.9% | 75.1% | ... | ... | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV COMORBIDITIES

| | |
|---|---------------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 3100 [2500–3600] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | 8.4% |
| Cervical cancer screening of women living with HIV | ... |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment | ... |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment | ... |

HIV TESTING AND TREATMENT CASCADE



| | ... | 64% | ... |
|-----------------|------------|----------|------------|
| All ages | [...-...%] | [54–78%] | [...-...%] |
| Children (0–14) | [...-...%] | [39–64%] | [...-...%] |
| Women (15+) | [...-...%] | [63–90%] | [...-...%] |
| Men (15+) | [...-...%] | [44–65%] | [...-...%] |

| | |
|---|-----|
| Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? | Yes |
|---|-----|

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|--------------------|-----------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | 42% [33–52%] | 70% [55–86%] |
| Early infant diagnosis | ...% [...-...%] | 40% [32–50%] |

HIV PREVENTION

| | |
|---|----------------|
| Knowledge of HIV prevention among young people aged 15–24 years (2017) | |
| — Women | 38.3% |
| — Men | 36.2% |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) | |
| — Women | ... |
| — Men | ... |
| Women aged 15–49 years who have their demand for family planning satisfied by modern methods (2012) | 45.6% |
| Men aged 15–49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | ... |

Harm reduction

| | |
|---|-----|
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | ... |
| — Safe injection rooms available (2016) | ... |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|---------------------------|---------------------------|---------------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | 2400 [1900–2900] | 2000 [1600–2400] | 1900 [1500–2500] |
| New HIV infections (0–14) | <100 [<100– <200] | <100 [<100– <100] | <100 [<100– <100] |
| New HIV infections (women, 15+) | ... [...–...] | ... [...–...] | ... [...–...] |
| New HIV infections (men, 15+) | ... [...–...] | ... [...–...] | ... [...–...] |
| HIV incidence per 1000 population | 0.87 [0.71–1.06] | 0.72 [0.58–0.88] | 0.66 [0.52–0.86] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | 2500 [2100–3100] | 1700 [1400–2100] | 1500 [1100–1900] |
| AIDS-related deaths (0–14) | <200 [<100– <200] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (women, 15+) | ... [...–...] | ... [...–...] | ... [...–...] |
| AIDS-related deaths (men, 15+) | ... [...–...] | ... [...–...] | ... [...–...] |
| People living with HIV | | | |
| People living with HIV (all ages) | 35 000 [29 000–40 000] | 34 000 [27 000–38 000] | 34 000 [28 000–39 000] |
| People living with HIV (0–14) | 680 [590–780] | <500 [<500–550] | <500 [<500– <500] |
| People living with HIV (women, 15+) | ... [...–...] | ... [...–...] | ... [...–...] |
| People living with HIV (men, 15+) | ... [...–...] | ... [...–...] | ... [...–...] |

LAWS AND POLICIES

| | |
|---|---|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | No, but prosecutions exist based on general criminal laws |
| Criminalization of sex work | Selling and buying sexual services is criminalized |
| Criminalization of same-sex sexual acts | Yes, imprisonment (up to 14 years) |
| Drug use or possession for personal use is an offence | Drug use or consumption is a specific offence in law |
| Criminalization of transgender people | Neither criminalized nor prosecuted |
| Laws or policies restricting the entry, stay and residence of people living with HIV | No |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 16 years |
| Spousal consent for married women to access sexual and reproductive health services | No |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | No |

STIGMA AND DISCRIMINATION

| | |
|---|------------|
| Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV | 2017 71 |
| Percentage of people living with HIV denied health services because of their HIV status in the last 12 months | |
| Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent | |


VIOLENCE

| | |
|---|--------------|
| Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months | 2012 9.89 |
|---|--------------|


EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|-----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2012 | US\$ 6 955 818 | US\$ 3 115 403 | ... | ... | ... | US\$ 17 136 135 |


EPIDEMIC TRANSITION METRICS




Change in new HIV infections since 2010 = **-6%**



Change in AIDS-related deaths since 2010 = **-16%**



Incidence:prevalence ratio = **0.06**




KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | 18 696 | ... | ... | ... | ... |
| HIV prevalence | 2% | ... | ... | ... | ... |
| Know their HIV status | 65.2% | ... | ... | ... | ... |
| Antiretroviral therapy coverage | ... | ... | ... | ... | ... |
| Condom use | 95% | ... | ... | ... | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV COMORBIDITIES

| | |
|---|------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 55 [41–72] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | ... |
| Cervical cancer screening of women living with HIV | ... |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment | ... |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment | ... |

HIV TESTING AND TREATMENT CASCADE



| | | | |
|-----------------|-----------------|-----------------|-----------------|
| All ages | 75% [63–87%] | 34% [29–40%] | 17% [15–20%] |
| Children (0–14) | >95% [>95–>95%] | 60% [48–76%] | 14% [11–18%] |
| Women (15+) | ...% [...–...%] | ...% [...–...%] | ...% [...–...%] |
| Men (15+) | ...% [...–...%] | ...% [...–...%] | ...% [...–...%] |

Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? **No**

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|-----------------|-----------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | 82% [69–>95%] | >95% [85–>95%] |
| Early infant diagnosis | ...% [...–...%] | ...% [...–...%] |

HIV PREVENTION

| | |
|--|----------------|
| Knowledge of HIV prevention among young people aged 15–24 years | ... |
| — Women | ... |
| — Men | ... |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) | ... |
| — Women | ... |
| — Men | ... |
| Women aged 15–49 years who have their demand for family planning satisfied by modern methods | ... |
| Men aged 15–49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | ... |
| Harm reduction | ... |
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | No |
| — Safe injection rooms available (2016) | No |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|----------------------|----------------------|----------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | <500 [<200– <500] | <500 [<200– <500] | <500 [<500– <500] |
| New HIV infections (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| New HIV infections (women, 15+) | <100 [<100– <200] | <200 [<100– <200] | <200 [<200– <500] |
| New HIV infections (men, 15+) | <100 [<100– <200] | <200 [<100– <200] | <200 [<200– <500] |
| HIV incidence per 1000 population | 0.41 [0.31–0.62] | 0.46 [0.37–0.69] | 0.56 [0.42–0.84] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | <200 [<200– <500] | <200 [<200– <200] | <200 [<100– <200] |
| AIDS-related deaths (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (women, 15+) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (men, 15+) | <100 [<100– <200] | <100 [<100– <200] | <100 [<100– <100] |
| People living with HIV | | | |
| People living with HIV (all ages) | 3700 [3300–4400] | 3800 [3400–5000] | 4800 [4300–6300] |
| People living with HIV (0–14) | <100 [<100– <200] | <100 [<100– <200] | <100 [<100– <200] |
| People living with HIV (women, 15+) | 1600 [1400–1900] | 1700 [1500–2300] | 2300 [2000–3000] |
| People living with HIV (men, 15+) | 2100 [1800–2500] | 2000 [1700–2600] | 2400 [2100–3100] |

LAWS AND POLICIES

| | |
|---|--|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | Yes |
| Criminalization of sex work | Partial criminalization of sex work |
| Criminalization of same-sex sexual acts | Laws penalizing same-sex sexual acts have been decriminalized or never existed |
| Drug use or possession for personal use is an offence | Drug use or consumption is a specific offence in law |
| Criminalization of transgender people | Neither criminalized nor prosecuted |
| Laws or policies restricting the entry, stay and residence of people living with HIV | No |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 16 years |
| Spousal consent for married women to access sexual and reproductive health services | No |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | Yes |

STIGMA AND DISCRIMINATION

Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent





VIOLENCE

Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months

EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2011 | US\$ 81 800 | US\$ 2 249 605 | ... | US\$ 1 009 894 | US\$ 12 385 | US\$ 4 674 508 |

EPIDEMIC TRANSITION METRICS



Change in new HIV infections since 2010 = 30%

Change in AIDS-related deaths since 2010 = -32%

Incidence:prevalence ratio = 0.06


KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | ... | ... | ... | ... | ... |
| HIV prevalence | 3.3% | 5.8% | ... | ... | ... |
| Know their HIV status | ... | ... | ... | ... | ... |
| Antiretroviral therapy coverage | ... | ... | ... | ... | ... |
| Condom use | ... | 53.3% | ... | ... | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV COMORBIDITIES

| | |
|---|------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 36 [28–46] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | ... |
| Cervical cancer screening of women living with HIV | ... |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment | ... |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment | ... |

HIV TESTING AND TREATMENT CASCADE



| | | | |
|-----------------|---------------|---------------|--------------|
| All ages | 65% [57–85%] | 51% [45–67%] | 43% [38–56%] |
| Children (0–14) | 92% [77–>95%] | 85% [72–>95%] | 63% [53–94%] |
| Women (15+) | 68% [59–88%] | 51% [45–66%] | 44% [38–57%] |
| Men (15+) | 61% [53–79%] | 49% [42–64%] | 41% [36–54%] |

Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? No

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|---------------|---------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | 82% [70–>95%] | 76% [67–>95%] |
| Early infant diagnosis | 9% [7–11%] | 67% [52–76%] |

HIV PREVENTION

| | |
|--|----------------|
| Knowledge of HIV prevention among young people aged 15–24 years | ... |
| — Women | ... |
| — Men | ... |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) | ... |
| — Women | ... |
| — Men | ... |
| Women aged 15–49 years who have their demand for family planning satisfied by modern methods | ... |
| Men aged 15–49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | ... |
| Harm reduction | ... |
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | No |
| — Safe injection rooms available (2016) | No |

EPIDEMIC ESTIMATES

| | 2005 | 2010 | 2017 |
|-------------------------------------|-----------------------|-------------------------|-------------------------|
| New HIV infections | | | |
| New HIV infections (all ages) | 770 [690–870] | 520 [<500–580] | <500 [<500– <500] |
| New HIV infections (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| New HIV infections (women, 15+) | <500 [<200– <500] | <200 [<200– <200] | <100 [<100– <200] |
| New HIV infections (men, 15+) | 530 [<500–610] | <500 [<500– <500] | <500 [<200– <500] |
| HIV incidence per 1000 population | 0.59 [0.53–0.67] | 0.39 [0.33–0.44] | 0.24 [0.17–0.29] |
| AIDS-related deaths | | | |
| AIDS-related deaths (all ages) | <500 [<500– <500] | <500 [<500– <500] | <500 [<500– <500] |
| AIDS-related deaths (0–14) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (women, 15+) | <100 [<100– <100] | <100 [<100– <100] | <100 [<100– <100] |
| AIDS-related deaths (men, 15+) | <500 [<500– <500] | <500 [<500– <500] | <500 [<200– <500] |
| People living with HIV | | | |
| People living with HIV (all ages) | 9500 [8500–10 000] | 11 000 [9500–12 000] | 11 000 [9600–12 000] |
| People living with HIV (0–14) | <200 [<200– <200] | <200 [<200– <200] | <200 [<200– <200] |
| People living with HIV (women, 15+) | 2900 [2600–3200] | 3400 [3100–3800] | 3900 [3500–4300] |
| People living with HIV (men, 15+) | 6500 [5600–7200] | 7000 [6100–7900] | 6800 [5800–7600] |

LAWS AND POLICIES

| | |
|---|---|
| Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission | ... |
| Criminalization of sex work | Selling sexual services is criminalized |
| Criminalization of same-sex sexual acts | Yes, imprisonment (14 years - life) |
| Drug use or possession for personal use is an offence | Possession of drugs for personal use is specified as a criminal offence |
| Criminalization of transgender people | ... |
| Laws or policies restricting the entry, stay and residence of people living with HIV | ... |
| Parental consent for adolescents to access HIV testing | Yes, for adolescents younger than 14 years |
| Spousal consent for married women to access sexual and reproductive health services | ... |
| Mandatory HIV testing for marriage, work or residence permits or for certain groups | ... |

STIGMA AND DISCRIMINATION

Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent


VIOLENCE

Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months


EXPENDITURES

| | Financing sources | | | | | Total |
|-----------------------------|-------------------|-----------------|-----------------------|----------------------------|---------------------------|-----------------|
| | Domestic private | Domestic public | International: PEPFAR | International: Global Fund | International: all others | |
| Last available report: 2009 | US\$ 277 991 | US\$ 11 415 268 | ... | ... | US\$ 1 408 776 | US\$ 13 810 965 |


EPIDEMIC TRANSITION METRICS




Change in new HIV infections since 2010 = **-37%**



Change in AIDS-related deaths since 2010 = **-21%**




Incidence:prevalence ratio = **0.03**



KEY POPULATIONS

| | Sex workers | Gay men and other men who have sex with men | People who inject drugs | Transgender people | Prisoners |
|---|-------------|---|-------------------------|--------------------|-----------|
| Estimated size of population | ... | ... | ... | ... | ... |
| HIV prevalence | ... | 31.6% | ... | ... | ... |
| Know their HIV status | ... | ... | ... | ... | ... |
| Antiretroviral therapy coverage | ... | ... | ... | ... | ... |
| Condom use | ... | 51% | ... | ... | ... |
| Coverage of HIV prevention programmes | ... | ... | ... | ... | ... |
| Avoidance of health care because of stigma and discrimination | ... | ... | ... | ... | ... |

HIV TESTING AND TREATMENT CASCADE



| | ... | 62% | ... |
|-----------------|------------|------------|------------|
| All ages | [...-...%] | [56-69%] | [...-...%] |
| Children (0-14) | [...-...%] | [71- >95%] | [...-...%] |
| Women (15+) | [...-...%] | [76-93%] | [...-...%] |
| Men (15+) | [...-...%] | [43-56%] | [...-...%] |

Is antiretroviral therapy provided in community settings (such as outside health facilities) for people who are stable on antiretroviral therapy? ...

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

| | 2010 | 2017 |
|---|--------------------|-------------------|
| Percentage of pregnant women living with HIV accessing antiretroviral medicines | 92% [81- >95%] | 81% [71-89%] |
| Early infant diagnosis | ...% [...-...%] | 94% [86- >95%] |

HIV COMORBIDITIES


| | |
|---|---------------|
| Estimated number of incident tuberculosis cases among people living with HIV (2016) | 47 [30-67] |
| Proportion of people living with HIV newly enrolled in HIV care with active tuberculosis (2016) | 5.3% |
| Cervical cancer screening of women living with HIV | ... |
| Proportion of people coinfecting with HIV and hepatitis B virus receiving combined treatment | ... |
| Proportion of people coinfecting with HIV and hepatitis C virus starting hepatitis C treatment | ... |

HIV PREVENTION

| | |
|--|----------------|
| Knowledge of HIV prevention among young people aged 15-24 years | |
| — Women | ... |
| — Men | ... |
| Condom use at last higher-risk sex (with a non-marital, non-cohabiting partner) | |
| — Women | ... |
| — Men | ... |
| Women aged 15-49 years who have their demand for family planning satisfied by modern methods | ... |
| Men aged 15-49 years who are circumcised | Not applicable |
| Male circumcisions performed according to national standards | Not applicable |
| People who received PrEP at least once during the reporting period (2017) | ... |

Harm reduction

| | |
|---|-----|
| — Use of sterile injecting equipment at last injection | ... |
| — Needles and syringes distributed per person who injects | ... |
| — Coverage of opioid substitution therapy | ... |
| — Naloxone available (2016) | ... |
| — Safe injection rooms available (2016) | ... |



METHODS

Methods for deriving UNAIDS estimates

INTRODUCTION

UNAIDS annually provides revised global, regional and country-specific modelled estimates using the best available epidemiological and programmatic data to track the HIV epidemic. Modelled estimates are required because it is impossible to count the exact number of people living with HIV, people who are newly infected with HIV or people who have died from AIDS-related illness in any country: doing so would require regularly testing every person for HIV and investigating all deaths, which is logistically impossible and ethically problematic. Modelled estimates—and the lower and upper bounds around these estimates—provide a scientifically appropriate way of describing HIV epidemic levels and trends.

PARTNERSHIPS IN DEVELOPING METHODS FOR UNAIDS ESTIMATES

Country teams use UNAIDS-supported software to develop estimates annually. The country teams are primarily comprised of demographers, epidemiologists, monitoring and evaluation specialists, and technical partners.

The software used to produce the estimates is Spectrum, which is developed by Avenir Health, and the Estimates and Projections Package, which is developed by the East–West Center.¹ The UNAIDS Reference Group on Estimates, Modelling and Projections provides technical guidance on the development of the HIV component of the software.²

A BRIEF DESCRIPTION OF METHODS USED BY UNAIDS TO CREATE ESTIMATES

For countries where HIV transmission is high enough to sustain an epidemic in the general population, available epidemiological data typically consist of HIV prevalence results from pregnant women attending antenatal clinics and from nationally representative population-based surveys. Many countries have historically conducted HIV sentinel surveillance among women attending antenatal clinics, which requires collecting data from a selection of clinics for several months every few years. More recently, many countries have stopped conducting sentinel surveillance and are now using the data from

the routine HIV tests conducted when pregnant women at antenatal clinics are tested as part of programmes for the prevention of mother-to-child transmission. These data avoid the need to conduct a separate surveillance effort, and they provide a complete set of data from all clinics instead of samples from specific sites.

The prevalence trends among pregnant women at antenatal clinics, whether determined from surveillance or routine data, can be used to inform estimates of national prevalence trends, whereas data from population-based surveys—which are conducted less frequently but have broader geographical coverage and also include men—are more useful for informing estimates of national HIV prevalence levels. Data from these surveys also contribute to estimating age- and sex-specific HIV prevalence levels and trends. For a few countries in sub-Saharan Africa that have not conducted population-based surveys, HIV prevalence levels are adjusted based on comparisons of antenatal clinic surveillance and population-based survey data from other countries in the region. HIV prevalence trends and numbers of people on antiretroviral therapy are then used to derive an estimate of HIV incidence trends.

Historically, countries with high HIV transmission have produced separate HIV prevalence and incidence trends for rural and urban areas when there are well-established geographical differences in prevalence. To better describe and account for further geographical heterogeneity, an increasing number of countries have produced subnational estimates (e.g. at the level of the province or state) that, in some cases, also account for rural and urban differences. These subnational or rural–urban estimates and trends are then aggregated to obtain national estimates.

In the remaining countries, where HIV transmission largely occurs among key populations at higher risk of HIV and the epidemic can be described as low-level, the estimates are derived from either surveillance among key populations and the general low-risk population, or from HIV case reporting data, depending on which data are most reliable in a particular country. In countries with high-quality HIV surveillance data among the key populations, the data from repeated HIV prevalence studies focused on key populations are used to derive

¹ More information on Avenir Health can be found at www.avenirhealth.org. The East–West Center website can be found at www.eastwestcenter.org.

² For more on the UNAIDS Reference Group on Estimates, Modelling and Projections, please visit www.epidem.org.

national estimates and trends. Estimates of the size of key populations are increasingly derived empirically in each country; when studies are not available, they are derived based on regional values and consensus among experts. Other data sources—including HIV case reporting data, population-based surveys and surveillance among pregnant women—are used to estimate the HIV prevalence in the general low-risk population. The HIV prevalence curves and numbers of people on antiretroviral therapy are then used to derive national HIV incidence trends.

For most countries in western and central Europe and North America—and many countries in Latin America, the Caribbean and the Middle East and North Africa that have insufficient HIV surveillance or survey data, but which have robust disease reporting systems—HIV case reporting and AIDS-related mortality data from vital registration systems are used directly to inform trends and levels in national HIV prevalence and incidence. These methods also allow countries to take into account evidence of underreporting or reporting delays in HIV case report data, as well as the misclassification of deaths from AIDS-related illness.

In all countries where UNAIDS supports the development of estimates, assumptions about the effectiveness of HIV programme scale-up and patterns of HIV transmission and disease progression are used to obtain age- and sex-specific estimates of (a) people living with HIV, (b) people newly infected with HIV, (c) people dying from AIDS-related illness and (d) other important indicators (including treatment programme coverage statistics). These assumptions are based on systematic literature reviews and analyses of research study data by scientific experts. Demographic population data, including fertility estimates, are derived from the United Nations Population Division's World Population Prospects 2017 data.

Selected inputs into the model—including the number of people on antiretroviral therapy and the number of women accessing services for the prevention of mother-to-child transmission of HIV by type of regimen—are reviewed and validated in partnership with the United Nations Children's Fund (UNICEF), the World Health Organization (WHO), the Global Fund to Fight AIDS, Tuberculosis and Malaria (the Global Fund), and selected technical partners.

Final country-submitted files containing the modelled outputs are reviewed at UNAIDS to ensure that the results are comparable across regions and countries and over time.

UNCERTAINTY BOUNDS AROUND UNAIDS ESTIMATES

The estimation software calculates uncertainty bounds around each estimate. These bounds define the range within which the true value lies (if it can be measured). Narrow bounds indicate that an estimate is precise, while wide bounds indicate greater uncertainty regarding the estimate.

In countries using HIV surveillance data, the quantity and source of the data available partly determine the precision of the estimates: countries with more HIV surveillance data have smaller ranges than countries with less surveillance data or smaller sample sizes. Countries in which a national population-based survey has been conducted generally have smaller ranges around estimates than countries where such surveys have not been conducted, while countries producing subnational estimates at the provincial level have wider ranges. In countries using HIV case reporting and AIDS-related mortality data, the number of years of data and the magnitude of the cases reported or the deaths from AIDS-related illness observed will contribute to the precision of the estimate.

The assumptions required to arrive at the estimate also contribute to the width of the ranges around the estimates: in brief, the more assumptions that are made, the wider the uncertainty range, since each assumption introduces additional uncertainties. For example, the ranges around the estimates of adult HIV prevalence are smaller than those around the estimates of HIV incidence among children, which require additional data on prevalence among pregnant women and the probability of mother-to-child HIV transmission, each of which have their own additional uncertainty.

UNAIDS is confident that the actual numbers of people living with HIV, people who are newly infected with HIV or people who have died from AIDS-related illness lie within the reported ranges. Over time, more and better data from countries will steadily reduce uncertainty.

IMPROVEMENTS TO THE 2018 UNAIDS ESTIMATES MODEL

Country teams create new Spectrum files every year. The files may differ from one year to the next for two reasons. First, new surveillance and programme data are entered into the model; this can change HIV prevalence and incidence trends over time, including for past years.

Second, improvements are incorporated into the model based on the latest available science and statistical methods that lead to the creation of more accurate trends in HIV incidence. Due to these improvements to the model and the addition of new data to create the estimates, the results from previous years cannot be compared with the results from this year. However, a full historical set of estimates are created each year, enabling a description of trends over time.

Between the previous estimates and the 2018 estimates, the following changes were applied to the model under the guidance of the UNAIDS Reference Group on Estimates, Modelling and Projections and based on the latest scientific evidence.

- Demographic data in the models were updated from the World Population Prospects 2015 estimates to the 2017 estimates.
- Assumptions about retention on antiretroviral therapy among pregnant women living with HIV were included.
- Aggregate routine data on prevalence among women attending antenatal clinics are now used to estimate the number of women living with HIV who are giving birth.
- Assumptions about the trends in HIV prevalence among pregnant women versus trends among the general population were updated.
- Annual HIV mortality probabilities among people on treatment in western and central Europe and North America were revised based on a special analysis conducted by the Antiretroviral Therapy Cohort Collaboration.
- An option was added in the model to prioritize allocation of treatment to individuals with the lowest CD4 count who had not yet initiated treatment.
- A new approach to fitting more complex incidence patterns for countries using case reporting and vital registration data is available.
- New methods to estimate the proportion of people dying before diagnosis and time from infection to diagnosis were incorporated into the model for countries using case reporting data to estimate incidence.

More detailed information on revisions to the 2018 model and Spectrum generally can be found at www.epidem.org.

MEASURING ANTIRETROVIRAL THERAPY COVERAGE

Since 2013, UNAIDS has provided the number and estimates of the proportion of all adults and children living with HIV who are on antiretroviral therapy (as opposed to those eligible for therapy according to national or international guidelines). This approach to estimating coverage reflects the WHO recommendations of starting antiretroviral therapy among everyone diagnosed as HIV-positive.

Countries report the number of people on treatment through the Global AIDS Monitoring (GAM) tool and Spectrum. Although those values come through routine data, they are likely to have some level of uncertainty if the country cannot deduplicate individuals who might receive medication from two different clinics or if there are delays in reporting data. Using results from data quality reviews through 2016, an estimated uncertainty—0.88 and 1.04 for the lower and upper bounds, respectively—was added to the number of people on treatment at the regional and global levels.

PUBLICATION OF COUNTRY-SPECIFIC ESTIMATES

UNAIDS aims to publish estimates for all countries with populations of 250 000 or more in 2017. For countries with populations of 250 000 or more that did not submit estimates, UNAIDS developed estimates using the Spectrum software that were based on published or otherwise available information. These estimates contributed to regional and global totals but were not published as country-specific estimates.

In countries with low-level epidemics, the number of pregnant women living with HIV is difficult to estimate. Many women living with HIV in these countries are sex workers or people who use drugs—or they are the sexual partners of gay men and other men who have sex with men or people who use drugs—making them likely to have different fertility levels than the general population. UNAIDS does not present estimates of mother-to-child HIV transmission, including estimates related to children in some countries that have concentrated epidemics, unless adequate data are available to validate these estimates. UNAIDS also does not publish these estimates for countries where the estimated number of pregnant women living with HIV is less than 100.

With regard to reporting incidence trends, if there are not enough historical data to state with confidence

whether a decline in incidence has occurred, UNAIDS does not publish data other than that from the most recent year; this prevents users from making inaccurate inferences about trends. Specifically, incidence trends are not published if there are fewer than four data points for the key population or if there have been no data for the past four years for countries using repeated survey or routine testing data. Trends prior to 2000 are not published for countries using case surveillance models if there is no early case surveillance or mortality data available.

Finally, UNAIDS does not publish country estimates when further data or analyses are needed to produce valid estimates. More information on the UNAIDS estimates and the individual Spectrum files for most countries can be found on the UNAIDS website (www.unaids.org). Resulting estimates can be found in the Aidsinfo section of the UNAIDS website (<http://aidsinfo.unaids.org/>).

Methods for deriving the 90–90–90 targets

INTRODUCTION

Starting in 2016, UNAIDS has provided estimates of global, regional and country-specific progress against the 90–90–90 targets. Progress towards these targets is directly monitored using three basic indicators:

- Indicator 1 (the first 90): the percentage of all people living with HIV who know their HIV status.
- Indicator 2 (the second 90): the percentage of people who know their HIV-positive status and are accessing treatment.
- Indicator 3 (the third 90): the percentage of people on treatment who have suppressed viral loads.

Metrics related to Indicators 2 and 3 can also be expressed as a percentage of all people living with HIV. When numbers or coverage of the treatment target are expressed relative to the total number of people living with HIV, this is called the “HIV testing and treatment cascade.” Using this approach, the second and third targets of the 90–90–90 targets translate into 81% coverage of antiretroviral therapy and 73% of people achieving viral suppression by 2020.

UNAIDS published its first set of global and regional testing and treatment cascades in 2015. Estimates of antiretroviral therapy coverage among people living with HIV are available going back to when treatment was first introduced. Results presented in this report supersede the previously published 2015 and 2016 values.

Since 2015, UNAIDS has also tracked progress towards the 90–90–90 targets by monitoring viral load testing access among people on treatment. If most people in the country are receiving a viral load test annually, as

recommended by WHO, we can have confidence in the accuracy of the estimate of viral suppression among all people living with HIV.

METHODS FOR MEASURING THE 90–90–90 TARGETS

To describe country-level progress against the 90–90–90 targets, UNAIDS analysed data on the number of people who knew their HIV status, the number of people on treatment and the number of people who were virally suppressed among those tested, as reported through the GAM tool and Spectrum.

A description of the GAM system and the treatment target-related indicators that countries report against are provided in the UNAIDS GAM 2018 guidelines (1). All programme data submitted to UNAIDS—including the number of people reported to know their status, the number of people accessing treatment and the number of people on treatment who are virally suppressed—were validated by UNAIDS and its partners prior to publication.

Country-submitted data that did not meet the required validation checks for quality either at the indicator level or across the treatment cascade were not published. Not all countries were able to report against all three prongs of the 90–90–90 targets.

The final set of country measures of progress against the 90–90–90 targets for 2015 through 2017 are available at <http://aidsinfo.unaids.org>. Complete treatment cascades were available for 53 countries in 2017. Upper and lower ranges of uncertainty for country-level estimates were calculated from the range of estimated numbers of people living with HIV. This range may not fully capture uncertainty in the reported programme data.

To estimate regional and global progress against the 90–90–90 targets, UNAIDS supplemented the country-supplied data submitted through GAM with data obtained from a review of other published and unpublished data sources, including grey literature and Demographic and Health Survey results. There were insufficient reported data from countries in western and central Europe and North America in 2017 to present results for the region, although the country values that were available in the region were used to construct the global totals. Upper and lower ranges of uncertainty for global and regional estimates were calculated from the range of numbers of people living with HIV and the lower and upper ranges of the numbers of people on treatment in the region. This range may not fully capture uncertainty in the reported or missing programme data for the first and third indicators.

DATA SOURCES AND INDICATOR-SPECIFIC METHODS FOR DERIVING GLOBAL AND REGIONAL METHODS

Estimates of people living with HIV

Unless otherwise stated, all progress measures in this report are based on UNAIDS global, regional and country-specific modelled estimates of the numbers of people living with HIV from Spectrum. Estimates of people living with HIV were available for 169 countries. More details about how UNAIDS derives estimates and uncertainty bounds around the number of people living with HIV and those accessing antiretroviral therapy can be found under “Measuring antiretroviral therapy coverage” (above, in Part 1 of this annex).

Knowledge of HIV status among people living with HIV

Global and regional measures of the number of people living with HIV who know their status were derived using the most recent HIV surveillance, programme data, nationally representative population-based survey data and modelled estimates for 102 countries in 2017. Where data were available separately for children (aged 0–14 years) and adults (aged 15 years and older), age-specific measures were first calculated and then aggregated to produce a national measure.

For 80 countries in 2017, the number of people living with HIV who knew their HIV status is based on HIV surveillance systems, programme registers or modelled estimates derived from case surveillance and programme data. If the measure from these sources was lower than the number of people accessing antiretroviral

therapy, the reported value was excluded from the analysis and replaced by a regionally-derived estimate. For countries using HIV surveillance or programme data, a country’s measure was included only if the HIV surveillance system had been functioning since before 2008. Countries with more recent systems may not have captured all people living with HIV who were diagnosed prior to 2008.

Although HIV surveillance systems, including those based on programme registers, can be a reasonably robust source of data to estimate the number of people living with HIV who know their status, biases in the reported numbers may still exist. For example, a country’s measure of the knowledge of status may be underestimated if not all people diagnosed are reported to the surveillance system in a timely manner; the measure also may be overestimated if people are reported to the system or included on a register more than once and these duplicates are not detected. Similarly, if people die or emigrate but are not removed from the system, the number of people living with HIV who are reported to know their HIV status also will be overstated.

The estimated numbers of people living with HIV who knew their status for 14 countries in sub-Saharan Africa in 2017 were derived from nationally representative population-based surveys conducted since 2011 and from treatment data reported through GAM. Four countries with surveys through 2017 directly asked respondents who tested HIV-positive whether they knew their HIV status as part of the survey, and this proportion was applied to the total number of people estimated to be living with HIV in the country. In the remaining 10 countries with a survey that did not directly ask participants about knowledge of their HIV status, a stepwise approach was used to estimate knowledge of status.

- In the first step, the total percentage of people who could know their status in the year of the most recent survey is estimated. For adults, this percentage is estimated by calculating the percentage of those who tested HIV-positive in the survey who had reported ever having been tested for HIV and had received the last test result. For children, who are not included in the survey, a proxy measure of treatment coverage in the survey year is used to estimate knowledge of status among children. This is a conservative measure, as some children may not have initiated treatment. To estimate knowledge of status for all people in the

year of the survey, the child and adult estimates are combined, weighted by the numbers of children and adults living with HIV.

- In the second step, the percentage of people who could know their status in the current or previous reporting year is derived by projecting the results from the first step forward. To do this, an assumption is made that the rate of testing scale-up in the era of test-and-treat was the same as the rate of scale-up of people starting treatment, calculated by the percentage point difference in total treatment coverage (for both adults and children) between the survey year and the treatment coverage value for either the current or previous year. For surveys conducted in 2017, the 2015 and 2016 values are estimated for previous years using a similar process as the one described above.
- In the third step, the estimate of people living with HIV who know their status for the year is derived by using the midpoint between the percentage of people living with HIV who could know their status (i.e. the second step) and the percentage of people living with HIV on treatment.

The measurement of knowledge of HIV status based on survey data when participants are not directly asked if they know their HIV status has several limitations. Typically, estimates derived from these surveys will underestimate knowledge of status for three reasons:

1. In settings where stigma and discrimination is or has been high, people may be reluctant to disclose that they have ever tested for HIV and received their results.
2. People who report ever testing may have seroconverted after their last test result and are therefore incorrectly counted as aware of their HIV status.
3. Most surveys that do not directly ask respondents about their HIV status occurred prior to 2017. Although surveys conducted prior to 2011 were excluded, it is possible that the adjustment method based on treatment scale-up does not accurately capture increases in the knowledge of status that occur over time among people living with HIV.

Underestimation of the reported number of people living with HIV who know their status can also occur in countries where survey respondents are directly asked

about their HIV status. In these instances, the risk is that survey participants do not disclose their HIV status to interviewers and are incorrectly classified as unaware of it. While it is impossible to measure the exact magnitude of this bias, in previous surveys in Kenya, Malawi and Uganda, anywhere from one tenth to one third of HIV-positive participants misreported their HIV status as negative (2). Underestimation of knowledge of status also can occur at the national level if people living with HIV learn their status either as a result of—or subsequent to—the survey, although this proportion of the total number of people in a country who know their status will be small.

For 34 countries without a current measure of knowledge of status in 2017, UNAIDS used published and unpublished grey literature and historical estimates reported through GAM to inform the regional and global values. A similar method used to project estimated knowledge of status for direct surveys from historical data was applied to estimates from such countries before 2017.

For 40 countries without any estimate of the number of people living with HIV who know their status—countries that are home to just 8% of the total estimated number of people living with HIV worldwide—the regional average of the ratio of the number of people who know their status and the number on treatment was calculated from available data submitted by countries in the region and weighted according to the number of people living with HIV by country. Knowledge of status was capped at 95%. The total number of people estimated to know their HIV status in countries was added across the region and globally to construct the numerator of the first 90 and the denominator of the second 90.

People accessing antiretroviral therapy

Global and regional measures of antiretroviral therapy numbers are calculated from country-reported programme data through GAM and the UNAIDS-supported Spectrum software. For a small number of countries where reported numbers of people on treatment are not available—primarily in western and central Europe and North America—estimates of the number of people on treatment are developed either in consultation with the public health agency responsible for monitoring the national treatment programme or based on published sources.

In partnership with UNICEF, WHO and other partners that support treatment service delivery in countries,

UNAIDS reviews and validates treatment numbers reported through GAM and Spectrum on an annual basis. UNAIDS staff also provide technical assistance and training to country public health and clinical officers to ensure the quality of the treatment data that are reported. Nevertheless, this measure may overestimate the number of people on treatment if people who transfer from one facility to another are reported by both facilities. Similarly, coverage may be overestimated if people who have died, disengaged from care or emigrated are not identified and removed from treatment registries. Treatment numbers also may be underestimated if not all clinics report the numbers on treatment completely or in a timely manner.

In 2016, UNAIDS completed a triangulation of data to verify the UNAIDS global estimate of people accessing antiretroviral therapy at the end of 2015. In 2018, UNAIDS has partnered with WHO, the Global Fund, selected technical partners and ministries of health in 28 countries (most in sub-Saharan Africa) to conduct data quality reviews of reported treatment numbers. For more details about how confident UNAIDS is in reported treatment numbers, please see *How many people living with HIV access treatment?*³

People who have achieved viral suppression

Progress towards the viral suppression target among people on treatment and as a proportion of all people living with HIV is derived from data reported to GAM. For the purposes of reporting, the threshold for suppression is a viral load of less than 1000 copies per ml, although some countries may set lower thresholds or require persons to achieve an undetectable viral load. This guidance also specifies that only a person's last test result from the reporting year be submitted, so the reported number suppressed among those tested should represent people and not tests performed.

UNAIDS GAM 2018 guidelines were updated from those of 2017 to include a threshold for reporting viral load suppression outcomes, such that testing coverage should be accessible to all or nearly all (>90%), or that it is nationally representative of people on treatment (typically 50–90% testing coverage). For countries with nationally representative but not universally accessible access to treatment, the estimate of viral suppression among those tested (i.e. the third 90) was multiplied by the number of people on treatment nationally to obtain overall viral suppression levels in the country.

Based on the more stringent coverage threshold, 67 countries reported viral load suppression data from case-based surveillance or laboratory-based reporting systems in 2018 (compared with 88 in 2017). Five countries had estimates based on nationally representative population-based surveys, where viral load testing was done only among those who self-reported that they were on treatment.

Estimates for the remaining countries were constructed using the regional average of the number of people on antiretroviral therapy who are virally suppressed, weighted according to the number of people on treatment in a country. The total number of people suppressed was added across the region and globally to construct the third 90 and the overall estimate of viral suppression among people living with HIV. The same approach also was used to construct historical regional and global estimates.

A number of challenges exist in using country-reported data to monitor the viral load suppression target.

- Routine viral load testing may not be offered at all treatment facilities, and those facilities where it is offered may not be representative of the care available at facilities without viral load testing. By assuming that the percentage of people suppressed among those accessing viral load testing is representative of all people on treatment in countries with incomplete viral load testing uptake, the measure may be either overestimated or underestimated depending on the characteristics of the reporting clinics where testing is available.
- Reported access to viral load testing varies considerably across each region, and it is difficult to know whether the experience in countries that reported data to UNAIDS is similar to that of countries in the same region that did not report data. In western and central Africa, for example, only 7 of 14 countries reported estimates of viral load suppression in 2017, representing just 14% of all people on treatment in the region. In Asia and the Pacific, nationally representative estimates of viral load suppression are not available for China and India in 2017. As a result, estimates for that region are constructed based on the remaining quarter of all people accessing treatment in the region where viral load suppression data are available.

³This document is available at <http://www.unaids.org/en/resources/documents/2016/how-many-people-living-with-HIV-access-treatment>.

- UNAIDS guidance requests routine (annual) viral load testing results only for people who are on treatment and eligible for testing. If people newly initiated on treatment achieve viral suppression but have not yet been offered viral load testing, they will be incorrectly classified as not suppressed and the resulting viral suppression estimate will be understated. UNAIDS also requests that countries only report results from routine viral load testing; if countries report test results that are primarily performed because of suspected treatment failure, the number of people virally suppressed in these countries will be underestimated. UNAIDS validates country submissions for quality, but it is not always possible to identify cases where both routine and other types of testing are occurring.
 - UNAIDS guidance recommends reporting viral load test results only for people on antiretroviral treatment; persons who naturally suppress the virus and are not on treatment will not be included in this measure.
- As access to viral load testing coverage expands and routine monitoring systems are strengthened to compile and report these data, the ability to quantify and eventually reduce bias in the 90–90–90 targets will improve.

Distribution of new HIV infections by subpopulation

The distribution of new HIV infections by region was estimated based on data for 169 countries using five data sources.

For countries that model their HIV epidemic based on data from subpopulations, including key populations, the numbers of new infections were extracted from Spectrum 2017 files. This source provided data for sex workers from 58 countries, for people who inject drugs from 36 countries, for gay men and other men who have sex with men from 56 countries, and for transgender people from 15 countries (all of which were located in Latin America, the Caribbean and Asia). Additionally, 21 countries (mostly from Asia) had data from clients of sex workers.

The second source was mode of transmission studies conducted in countries between 2006 and 2012. The proportions of new infections estimated for each subpopulation, calculated by modes of transmission analyses, were multiplied by the number of total new gender-specific adult infections (among those aged 15–49 years) to derive an estimated number of new infections by subpopulation. This source provided data for sex workers from 18 countries, for people who inject drugs from 25 countries, and for gay men and other men who have sex with men from 22 countries.

New HIV infections for European countries with neither of the aforementioned data sources were derived from

the European Centre for Disease Prevention and Control (ECDC) and World Health Organization Regional Office for Europe *HIV/AIDS surveillance in Europe 2017–2016 data* (3). The proportions of new diagnoses for each region in Europe (West, central and East) were applied to UNAIDS estimates of new infections in each country for people who inject drugs and gay men and other men who have sex with men. Data for sex workers were not available from the ECDC report. New HIV infections in China, the Russian Federation and the United States were taken from the most recent available national reports of new diagnoses.

New HIV infections among countries without a direct data source were calculated from regional benchmarks. The benchmarks were set by the median proportion of new infections in the specific subpopulation in all available countries in the same region. The majority of these countries were located in sub-Saharan Africa. There were 73 countries that used benchmark values for the sex work estimate, 95 countries for the people who inject drugs estimate, 33 countries for the gay men and other men who have sex with men estimate, and 36 countries for the transgender people estimate.

The calculated proportions of infections for each key population include the sex partners of members of key populations. New infections among sex partners of key populations were estimated using the number of sex partners and transmission probabilities from the literature.

References

1. Global AIDS monitoring 2018: indicators for monitoring the 2016 United Nations Political Declaration on HIV and AIDS. Geneva: UNAIDS; 2018 (http://www.unaids.org/sites/default/files/media_asset/2017-Global-AIDS-Monitoring_en.pdf).
2. Johnston LG, Sabin ML, Prybylski D, Sabin K, McFarland W, Baral S et al. Policy and practice: the importance of assessing self-reported HIV status in bio-behavioural surveys. *Bull World Health Organ.* 2016;94:605–12.
3. European Centre for Disease Prevention and Control (ECDC), World Health Organization Regional Office for Europe. HIV/AIDS surveillance in Europe 2017–2016 data. Stockholm: ECDC; 2017.

Copyright: © 2018

Joint United Nations Programme on HIV/AIDS (UNAIDS)

All rights reserved.

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of UNAIDS concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. UNAIDS does not warrant that the information published in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use.

UNAIDS/JC2924/Caribbean



CARIBBEAN

UNAIDS
Joint United Nations
Programme on HIV/AIDS

20 Avenue Appia
1211 Geneva 27
Switzerland

+41 22 791 3666

unaids.org