
Figure 1.1

The virtual elimination of mother-to-child transmission of HIV is possible

Estimated New HIV infections among children 0-14: Different scenarios for 25 countries



- No ARV prophylaxis for PMTCT
- Constant 2009 coverage of ARV prophylaxis
- 90% of women reached with services matching WHO guidelines
- 90% of women reached with services matching WHO guidelines, incidence reduced by 50%, and eliminate unmet need for family planning
- 90% of women reached with services matching WHO guidelines, incidence reduced by 50%, eliminate unmet need for family planning, restrict breastfeeding to 12 months



Figure 1.2

Millennium Development Goal 6 indicators

Population-adjusted averages for indicators for Millennium Development Goal target 6.A (halt and begin to reverse the spread of HIV/AIDS), 1999–2003 and 2004–2009.






Figure 1.3

Young people and sexual risk

People aged 15–25 years who had sex before age 15 years and who had multiple partners in the past 12 months.





Figure 1.4

Treatment coverage in low- and middle-income countries

Population-adjusted averages for treatment coverage in low- and middle-income countries by geographical region in 2009 based on 2010 WHO guidelines: Millennium Development Goal target 6.B (achieve, by 2010, universal access to treatment for HIV/AIDS for all those who need it). The regional figure for North America is not shown because of lack of data.

% 10 20 30 40 50 60

Sub-Saharan Africa
Latin America
Caribbean

East, South and South-east Asia
Europe and Central Asia
North Africa and the Middle East

