


GLOBAL REPORT


Figure 3.1 (1/2)

HIV prevention in Zambia, 2000-2009

**Males
15–24
years**



**Females
15–24
years**




- Young people never having had sex (15-19 years)
- Young people never having had sex (15-24 years)
- Never married respondents who had sex in the last 12 months who used a condom with their last sexual partner
- Condom use at last sex with a non-marital, non-cohabiting sexual partner
- Condom use at last sex, among respondents who had >1 partner in the last year
- Proportion of respondents who had had sex before age 15
- Proportion of respondents that had >1 partner in past year

Source: Zambia Sexual Behavior Survey.

GLOBAL REPORT

Figure 3.1 (2/2)

HIV prevention in Zambia, 2000-2009



Source: Zambia Sexual Behavior Survey.


GLOBAL REPORT

Figure 3.2

Young people's knowledge of HIV

Countries with comprehensive correct knowledge of HIV exceeding 60% among people 15–24 years old.

.....




Source: Country Progress Reports 2010.

GLOBAL REPORT

Figure 3.3

Multiple sexual partners in the past year, Uganda

Percentage of the population (ages 15–49 years old) that have had multiple sex partners in the past year in Uganda, by sex and age group, 1989–2006.



Source: Demographic and Health Surveys and other population-based behavioural survey data.

Association between harm reduction programmes and HIV prevalence in Ukraine, 2004–2009.

Source: M Mahy, C Chhea, T Saliuk, O Varetska, R Lyerla (2010). A proxy measure for HIV incidence among populations at increased risk to HIV Vol 2(1):8, Journal of HIV/AIDS Surveillance and Epidemiology. UNGASS Country Progress Reports 2010.


GLOBAL REPORT

Figure 3.5

Availability of sterile injecting equipment, 2010

Global estimates of the availability of sterile injecting equipment per person who uses drugs per year, 2010.

.....



Source: Mathers BM, Degenhardt L, Ali H, Wiessing L, Hickman M, Mattick R, et al. HIV prevention, treatment and care for people who inject drugs: A systematic review of global, regional and national coverage. The Lancet 2010;379:1014-28.

Table 3.1

Countries in which HIV infections among people who inject drugs represent 20% or more of the total number of people living with HIV

.....

Azerbaijan	Malaysia
Canada	New Zealand
China	Pakistan
Estonia	Russian Federation
Georgia	Spain
Indonesia	Tajikistan
Iran (Islamic Republic of)	Ukraine
Italy	United States of America
Kazakhstan	Uzbekistan
Kyrgyzstan	

Source: Mathers et al. (12).

GLOBAL REPORT

Figure 3.6

HIV prevention programmes for selected populations

Median coverage of HIV prevention programmes for selected population groups, 2008 and 2010.

.....




Source: Country Progress Reports 2010.

GLOBAL REPORT

Figure 3.7

Condom use by men who have sex with men

Percentage (median and range) of men who have sex with men who used a condom at last sex by geographical region, 2010.



Source: Country Progress Reports 2010.


GLOBAL REPORT

Figure 3.8

Condom use and HIV prevalence among sex workers in Cambodia

Percentage of sex workers using condoms and HIV prevalence among brothel-based sex workers in Cambodia by length of time involved in sex work, 1998–2007.

.....




Source: M Mahy, C Chhea, T Saliuk, O Varetska, R Lyerla (2010). A proxy measure for HIV incidence among populations at increased risk to HIV Vol 2(1):8, Journal of HIV/AIDS Surveillance and Epidemiology.

Figure 3.9

Averting HIV infection by eliminating police beatings of people who inject drugs, Ukraine

HIV infections that could be averted by eliminating police beatings of people who inject drugs in three cities in Ukraine

.....




Source: Strathdee et al 2010

Figure 3.10

Gaps in antiretroviral therapy to prevent mother-to-child transmission

Global gap in providing antiretroviral therapy to 80% of mothers to prevent mother-to-child transmission in low and middle-income countries.




Source: WHO Towards Universal Access 2010.

GLOBAL REPORT

Figure 3.11

Distribution of prophylactic regimens for the prevention of mother-to-child transmission



Source: Country Progress Reports 2010.

Coverage of antenatal care services and services for preventing mother-to-child transmission among women living with HIV in high-prevalence countries, 2010



UNAIDS
UNICEF
WFP
UNFPA
UNODC
ILO
UNESCO
WHO
WORLD BANK

GLOBAL REPORT

Table 3.2

Scaling up male circumcision

Recent roll-out of the scaling up of adult male circumcision in nine countries.

	Number circumcised	Time period	Number of sites established
BOTSWANA	6 180	April 2009 – March 2010	35
KENYA	91 300 (90 000 in Nyanza alone)	2009 – June 2010	
NAMIBIA	350	September 2009 – June 2010	3
RWANDA	542	October 2009 – April 2010	9
SWAZILAND	10 000	2008 – June 2010	
UGANDA	5 340	October 2008 – March 2010	
UNITED REPUBLIC OF TANZANIA	4 700	September 2009 – May 2010	3
ZAMBIA	9 906 10 000 9 179	January – June 2010 2009 2007 – 2008	56
ZIMBABWE	6 070	May 2009 – April 2010	5

Source: Meeting reports and presentations. Durham, NC, Clearinghouse on Male Circumcision for HIV Prevention, 2010.