

CAMEROON

Cameroon has shown a decline of 49% in the number of new HIV infections among children since 2009. In 2015 an estimated 82% of pregnant women living with HIV accessed antiretroviral medicines for the prevention of mother-to-child transmission. Challenges remain in maintaining women on antiretroviral medicines throughout the breastfeeding period, as the mother-to-child transmission rate of 5% at six weeks rises to 13% after breastfeeding ends. Paediatric diagnosis and treatment are areas for continued attention: in 2015, only 30% of infants exposed to HIV received early infant diagnosis of HIV, and 18% of children living with HIV received antiretroviral therapy. Cameroon is strengthening the capacity of home-based and community-based organizations to increase demand and support for prevention of mother-to-child transmission services.



4100

NEW INFECTIONS AMONG CHILDREN IN 2015



8 OUT OF 10

PREGNANT WOMEN LIVING WITH HIV ACCESSING ANTIRETROVIRAL MEDICINES TO PREVENT MOTHER-TO-CHILD-TRANSMISSION OF HIV



49%

DECREASE IN THE NUMBER OF NEW HIV INFECTIONS AMONG CHILDREN, 2009–2015




13%

RATE OF MOTHER-TO-CHILD HIV TRANSMISSION, INCLUDING DURING BREASTFEEDING


NEW HIV INFECTIONS AMONG CHILDREN

Number of new HIV infections among children (aged 0–14 years)




ART COVERAGE AMONG CHILDREN

Percentage of children (aged 0–14 years) living with HIV who have access to antiretroviral therapy




COVERAGE VS. TRANSMISSION

Increasing coverage of antiretroviral medicines has translated into decreasing rates of HIV transmission from mother to child



FAMILY PLANNING—UNMET NEED


Percentage of unmet need for family planning



Source: Demographic and Health Survey, 2011, all currently married women (aged 15–49 years).


EARLY INFANT DIAGNOSIS

Percentage of infants born to women living with HIV receiving a virological test by two months of age



WOMEN ACQUIRING HIV INFECTION

The number of women (aged 15–49 years) acquiring HIV increased by 5% since 2009



Source: UNAIDS 2016 estimates, unless otherwise noted. Every year, countries update their HIV estimates, including the historical trends of those estimates. The 2016 results reflect updated surveillance and programme data and improved models. The 2016 estimates are likely to differ to estimates produced in earlier years and are believed to be more accurate. Paediatric antiretroviral therapy coverage is based on all children living with HIV, while in previous years it was restricted to children eligible for antiretroviral therapy.