


TUBERCULOSIS AND HIV

IN 2022, AN ESTIMATED 10.6 MILLION PEOPLE FELL ILL WITH TB AND 1.3 MILLION DIED FROM THE DISEASE



People living with HIV are up to **16 [14–18] times** more likely to fall ill **with TB**

ANNUAL GLOBAL FUNDING FOR DEVELOPMENT AND RESEARCH ON TUBERCULOSIS IS **US\$ 1.0 BILLION** SHORT OF WHAT IS REQUIRED



TB IS THE LEADING CAUSE OF DEATH AMONG PEOPLE LIVING WITH HIV

UNAIDS IS WORKING WITH PARTNERS TO REDUCE TB-RELATED DEATHS AMONG PEOPLE LIVING WITH HIV BY **80%** (COMPARED TO A 2010 BASELINE) AND THAT **90%** OF PEOPLE LIVING WITH HIV RECEIVE PREVENTIVE TREATMENT FOR TUBERCULOSIS **BY 2025**



TB is curable:
the combination of TB treatment and ART is estimated to have averted 6.4 million deaths between 2010 and 2022

In 2022, only 54% of people living with HIV estimated to have developed TB received both TB treatment and ART

GLOBALLY, 11.3 MILLION PEOPLE LIVING WITH HIV RECEIVED TB PREVENTIVE TREATMENT BETWEEN 2018 AND 2022

IN 2022, AN ESTIMATED 170 000 PEOPLE DIED FROM AIDS-RELATED TB

SIMPLE, AFFORDABLE AND EFFECTIVE HIV/TB PROGRAMMES

All people living with HIV should have access to:

- Antiretroviral Therapy
- Regular TB screening
- TB diagnostics and treatment
- TB preventive therapy (if no symptoms)



All people living with TB should have access to:

- HIV testing and antiretroviral therapy
- HIV prevention options
- TB treatment